ලී ලංකා විතාශ දෙපාර්තමේන්තුව ලී ලංකා විතාශ දෙපාර්තමේන**ිලල්යුණුවේන්තුල පළුජාවේන්තුවා** විතාශ දෙපාර්තමේන්තුව ලී ලංකා විතාශ දෙපාර්තමේන්තුව இலங்கைப் பரீட்சைத் திணைக்களம் இலங்கைப் பரீட்சைத் திணைக்களம்இள்ளது பரீட்சைத் திணைக்களம் இலங்கைப் பரீட்சைத் திணைக்களம் Department of Examinations, Sri Lanka Department of **இතාස්තෙසට Sri Liftlik නෙම**නු tra<del>නි</del> කාත්ත සහාර්තමේන්තුව ලී ලංකා විතාශ දෙපාර්තමේන්තුව ලී ලංකා විතාශ සහාර්තමේන්තුව ලී ලංකා විතාශ දෙපාර්තමේන්තුව ලී ලංකා විතාශ සහාර්තමේන්තුව ලී ලංකා විතාශ සහාර්තමේන්තුව ලී ලංකා විතාශ දෙපාර්තමේන්තුව ලී ලංකා විතාශ සහාර්තමේන්තුව ලී ලංකා විතාශ දෙපාර්තමේන්තුව ලී ලංකා විතාශ දෙපාර්තමේන්තුව ලී ලංකා විතාශ දෙපාර්තමේන්තුව ලේ ලංකා විතාශ දෙපාර්තමේන්තුව ලේ ලංකා විතාශ දෙපාර්තමේන්තුව ලේ ලේකා විතාශ දෙපාර්තමේන්තුව ලේකා විතාශ දෙපාර දෙපාර

අධායන පොදු සහතික පතු (උසස් පෙළ) විභාගය, 2018 අශෝස්තු <u>கல்விப் பொதுத் தராதரப் பத்திர (உயர் தர)ப் பரீட்சை, 2018 ஓகஸ்ங்</u> General Certificate of Education (Adv. Level) Examination, August 2018

## 15.08.2018 / 0830 - 1030

රසායන විදහාව இரசாயனவியல் Chemistry



පැය ලදකයි இரண்டு மணித்தியாலம் Two hours

#### **Instructions:**

- \* Periodic Table is provided.
- This paper consists of 09 pages.
- Answer all the questions.
- Use of calculators is not allowed.
- Write your Index Number in the space provided in the answer sheet.
- Follow the instructions given on the back of the answer sheet carefully.
- In each of the questions 1 to 50, pick one of the alternatives from (1), (2), (3), (4), (5) which is correct or most appropriate and mark your response on the answer sheet with a cross (x) in accordance with the instructions given on the back of the answer sheet.

Universal gas constant  $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$ 

 $N_{\rm A} = 6.022 \times 10^{23} \, \rm mol^{-1}$ Avogadro constant

 $h = 6.626 \times 10^{-34} \text{ J s}$ Planck's constant

 $c = 3 \times 10^8 \text{ m s}^{-1}$ Velocity of light

- 1. The number of unpaired electrons present in a gaseous Co3+ ion in its ground state is,
  - (1) 1

- (4) 4
- 2. Which quantum number(s)  $(n, l, m_l, m_s)$  is/are associated with the shape of an atomic orbital of an atom?
  - (1) l
- (2)  $m_i$
- (3) n and l
- (4) n and  $m_i$
- (5) l and  $m_1$
- 3. What is the IUPAC name of the compound shown below?

$$\begin{array}{ccc} \operatorname{CH_3CH_2CH} & -\operatorname{C} = \operatorname{CHCO_2H} \\ & | & | \\ & \operatorname{Br} & \operatorname{NO_2} \end{array}$$

(1) 4-bromo-3-nitro-2-hexenoicacid

(2) 4-bromo-3-nitro-2-hexenoic acid

(3) 3-nitro-4-bromo-2-hexenoicacid

(4) 3-nitro-4-bromo-2-hexenoic acid

- (5) 3-bromo-4-nitro-4-hexenoic acid
- 4. The correct answer when the molecules  $O_2$ ,  $H_2O_2$ ,  $OF_2$  and  $O_2F_2$  (structure similar to  $H_2O_2$ ) are arranged in the decreasing order of the oxidation state of oxygen (O) is,
  - (1)  $O_2F_2 > OF_2 > O_2 > H_2O > H_2O_2$

(2)  $H_2O > H_2O_2 > O_2 > O_2F_2 > OF_2$ 

(3)  $H_2O_2 > O_2F_2 > O_2 > OF_2 > H_2O_2$ 

(4)  $OF_2 > O_2F_2 > O_2 > H_2O > H_2O_2$ 

- (5)  $OF_2 > O_2F_2 > O_2 > H_2O_2 > H_2O_3$
- 5. The most acceptable Lewis structure for the thiocyanate ion SCN is,
  - $(1) \quad \overset{\ominus}{:} \overset{\Box}{:} \overset{\Box}{:} \overset{\Box}{:} \overset{\ominus}{:} \overset{\Box}{:} \overset{\ominus}{:} \overset{\Box}{:} \overset{\ominus}{:} \overset{\Box}{:} \overset{\ominus}{:} \overset{\Box}{:} \overset{\Box}{:$

- 6. The molarity (mol dm<sup>-3</sup>) of a NaI solution which has a density of 1.03 g cm<sup>-3</sup> and is 3% NaI by mass is,

(Na = 23, I = 127)

- (1) 0.21
- (2) 0.23
- (3) 0.25
- (4) 0.28
- (5) 0.30

7. Precipitates of AgI and AgBr were added to a small amount of distilled water. This mixture was allowed to reach equilibrium at 25 °C. It was observed that both the solids were present in the system at equilibrium. Which of the following relations is applicable to this solution?

$$(K_{\rm sp(AgI)} = 8.0 \times 10^{-17} \, \text{mol}^2 \, \text{dm}^{-6}, K_{\rm sp(AgBr)} = 5.0 \times 10^{-13} \, \text{mol}^2 \, \text{dm}^{-6} \, \text{at 25 °C})$$

(1) 
$$[Br^-] = \sqrt{5.0 \times 10^{-13}} \mod dm^{-3}$$
 and  $[I^-] = \sqrt{8.0 \times 10^{-17}} \mod dm^{-3}$ 

(2) 
$$[Br^-][I^-] = [Ag^+]^2$$

(3) 
$$\left[Ag^{+}\right] = \left(\sqrt{5.0 \times 10^{-13}} + \sqrt{8.0 \times 10^{-17}}\right) \text{ mol dm}^{-3}$$

(4) 
$$\frac{[Br^-]}{[I^-]} = \frac{5.0}{8.0} \times 10^4$$

(5) 
$$[Ag^+] = [Br^-] = [I^-]$$

- 8. Which of the following statements is false?
  - (1) Although the carbonates of all the group two metals in the Periodic Table are insoluble in water, their bicarbonates are soluble.
  - (2) The hydroxides of all the group two metals in the Periodic Table are soluble in water.
  - (3) The nitrates of all the group two metals in the Periodic Table are soluble in water.
  - (4) The oxides and hydroxides of Na and Mg show basic properties whereas the oxide and hydroxide of Al show amphoteric properties.
  - (5) The hydrides of Si and S show weakly acidic properties.
- 9. In which list are the elements given in the order of increasing (left to right) atomic radii?
  - (1) Li, Na, Mg, S

(2) C, Si, S, Cl

(3) B, C, N, P

(4) Li, Na, K, Ca

- (5) B, Be, Na, K
- 10. Liquids A and B form an ideal solution. Consider a mixture of liquids A and B in equilibrium with the vapour in a closed rigid container at constant temperature.  $P_A^o$  and  $P_B^o$  respectively are the saturated vapour pressures of A and B while P is the total pressure of the container and  $X_A^g$  is the mole fraction of A in the vapour phase. Which of the following is correct about this system?

$$(1) \quad P = \left(P_A^o - P_B^o\right) X_A^g + P_B^o$$

(2) 
$$\frac{1}{P} = \left(\frac{1}{P_A^o} - \frac{1}{P_B^o}\right) X_A^g + \frac{1}{P_B^o}$$

(3) 
$$P = \left(P_A^o + P_B^o\right) X_A^g - P_B^o$$

(4) 
$$\frac{1}{P} = \left(\frac{1}{P_B^o} - \frac{1}{P_A^o}\right) \frac{1}{X_A^g}$$

(5) 
$$\frac{1}{P} = \left(\frac{1}{P_A^o} - \frac{1}{P_B^o}\right) \frac{1}{X_A^g}$$

11. The increasing order of boiling points of the following substances is,

He, CH<sub>4</sub>, CCl<sub>4</sub>, CBr<sub>4</sub>, SiH<sub>4</sub>

- (2) He < SiH<sub>4</sub> < CH<sub>4</sub> < CCl<sub>4</sub> < CBr<sub>4</sub>
- (1)  $CH_4 < He < SiH_4 < CCl_4 < CBr_4$ (3)  $He < CH_4 < SiH_4 < CCl_4 < CBr_4$
- (4)  $CH_4$  < He <  $SiH_4$  <  $CBr_4$  <  $CCl_4$
- (5) He < CH<sub>4</sub> < CCl<sub>4</sub> < SiH<sub>4</sub> < CBr<sub>4</sub>
- 12. Identify the correct statement from the following.
  - (1) Among the electronic transitions  $n=2 \longrightarrow n=1$ ,  $n=3 \longrightarrow n=2$  and  $n=4 \longrightarrow n=3$  in a hydrogen atom, most energy is released in  $n=3 \longrightarrow n=2$ .
  - (2) Among the species OF<sub>2</sub>, OF<sub>4</sub> and SF<sub>4</sub>, the least stable is SF<sub>4</sub>.
  - (3) Among the elements Li, C, N, Na and P, the least electronegative element is Li.
  - (4) In the following pairs (Li & F), (Li<sup>+</sup> & F<sup>-</sup>), (Li<sup>+</sup> & O<sup>2-</sup>) and (O<sup>2-</sup> & F<sup>-</sup>), the difference in radii is greatest between Li<sup>+</sup> and O<sup>2-</sup>.
  - (5) The only type of intermolecular force present in CH<sub>2</sub>Cl<sub>2</sub> in the liquid phase is dipole-dipole forces.

<b>13</b> .	Consider	the	reaction:	$CH_4(g) \longrightarrow$	$CH_3(g)$	+	H(g)
-------------	----------	-----	-----------	---------------------------	-----------	---	------

The standard change in enthalpy of the above reaction is,

- (1) the standard enthalpy change for the dissociation of the first C-H bond in methane.
- (2) the standard atomisation enthalpy change of methane.
- (3) the standard first ionisation enthalpy change of methane.
- (4) the standard bond dissociation enthalpy change of methane.
- (5) the standard radical formation enthalpy change of methane.
- 14. The elementary reaction  $2A(g) \longrightarrow B(g)$  occurs in a closed rigid container at a constant temperature. Initial pressure of the container is  $P_0$  and the pressure when the rate of reaction is 50% of the initial value is  $P_t$ . Which of the following gives the correct value for  $\frac{P_t}{P}$ ?

(1) 
$$\frac{P_t}{P_t} = \frac{1}{2}$$

(2) 
$$\frac{P_t}{P_0} = \frac{1}{\sqrt{2}}$$

(3) 
$$\frac{P_t}{P_o} = \frac{1+\sqrt{2}}{2\sqrt{2}}$$

values 4.7 and 5.0 respectively is at equilibrium. The value of  $\log \left(\frac{[A^-]}{[B^-]}\right)$  is approximately equal to,


$$(1) \quad \frac{P_t}{P_o} = \frac{1}{2} \qquad \qquad (2) \quad \frac{P_t}{P_o} = \frac{1}{\sqrt{2}} \qquad \qquad (3) \quad \frac{P_t}{P_o} = \frac{1+\sqrt{2}}{2\sqrt{2}} \quad (4) \quad \frac{P_t}{P_o} = \frac{\sqrt{2}}{1+\sqrt{2}} \quad (5) \quad \frac{P_t}{P_o} = \frac{\sqrt{2}-1}{1+\sqrt{2}}$$


- 15. An equimolar aqueous solution of the weak acids HA and HB (1.0 mol dm<sup>-3</sup> in each acid) with  $pK_{s}$ 
  - (1) 23.5
- (2) -0.3
- (3) 0.3
- (4) 0.94
- (5) 1.06
- 16. Which of the following statements about C<sub>6</sub>H<sub>5</sub>OH is false?
  - (1) Reacts with CH<sub>2</sub>COCl to form a phenyl ester.
  - (2) Reacts with bromine water to give a white precipitate.
  - (3) Evolves CO<sub>2</sub> gas when treated with NaHCO<sub>3</sub>.
  - (4) Gives a coloured compound when treated with  $C_6H_5N_2^+Cl^-$  in the presence of NaOH.
  - (5) Gives a coloured (purplish) solution when treated with neutral FeCl<sub>3</sub>.
- 17. The half life of a reaction is,
  - (1) always independent of the initial concentration of reactants.
  - (2) always dependent on the rate constant.
  - (3) always independent of the order of the reaction.
  - (4) always independent of temperature.
  - (5) equal to twice the total reaction time.
- 18. Electromotive force of an electrochemical cell does not depend on,
  - (1) the nature of the electrolytes.
  - (2) temperature.
  - (3) the concentrations of the electrolytes.
  - (4) the surface areas of the electrodes.
  - (5) the types of metals that form the electrodes.
- 19.  $IO_3^-$  (iodate ion) oxidizes the  $SO_3^{2-}$  ion to  $SO_4^{2-}$  in acidic medium. The mass of  $KIO_3$  required to totally oxidize the amount of Na<sub>2</sub>SO<sub>3</sub> present in 25.0 cm<sup>3</sup> of a solution of Na<sub>2</sub>SO<sub>3</sub> (0.50 mol dm<sup>-3</sup>) to  $Na_2SO_4$  is 1.07 g. (O = 16, K = 39, I = 127)

The final oxidation state of iodine after the completion of the reaction is,

- (1) -1
- (2) 0
- (3) +1
- (4) +2
- (5) +3
- 20. Which of the following statements is false with regard to the s-block elements in the Periodic Table?
  - (1) All elements in group I react with water liberating H<sub>2</sub> gas.
  - (2) All elements in group I except Li react with  $N_2$  gas.
  - (3) All elements in group II react with N<sub>2</sub> gas.
  - (4) Na reacts with excess  $O_2$  to give  $Na_2O_2$  whereas K gives  $KO_2$ .
  - (5) All elements in the s-block are good reducing agents.

21. A system consisting of two rigid containers containing an ideal gas is shown in the diagram. The containers can be connected to each other by opening the tap. The system changes from configuration A to configuration B when the tap is opened. In general n, P, V and T represent number of moles, pressure, volume and temperature respectively.





configuration A (tap closed)

configuration B (tap opened)

Which of the following relations is correct about this system?

$$(1) \ P_1 V_1 = P_2 V_2$$

(2) 
$$\frac{P_3 T_1}{P_1} + \frac{P_3 T_2}{P_2} = 2T_3$$

(3) 
$$\frac{T_1}{P_1} = \frac{T_2}{P_2}$$

(4) 
$$P_1T_1 = P_2T_2$$

(5) 
$$P_1V_1 + P_2V_2 = P_3(V_1 + V_2)$$

- 22. Which of the following statements is false with regard to 3d-elements of the Periodic Table?
  - (1) Atomic radii are smaller than the atomic radii of the s-block elements in the same period.
  - (2) Densities are higher than the densities of the s-block elements in the same period.
  - (3) V<sub>2</sub>O<sub>5</sub>, CrO<sub>3</sub> and Mn<sub>2</sub>O<sub>7</sub> are acidic oxides.
  - (4) First ionization energies are less than the first ionization energies of the s-block elements in the same period.
  - (5) The most common oxidation states of cobalt in cobalt compounds are +2 and +3.
- 23. Standard Gibbs energy changes for the reaction,  $MO(s) \rightarrow M(s) + \frac{1}{2}O_2(g)$  at two different temperatures are given below.

The standard entropy change of the reaction is,

- (1) 248.8 J K<sup>-1</sup> mol<sup>-1</sup>
- (2)  $-248.8 \text{ J K}^{-1} \text{ mol}^{-1}$ (5)  $48.4 \text{ J K}^{-1} \text{ mol}^{-1}$
- (3)  $-48.4 \text{ J K}^{-1} \text{ mol}^{-1}$

- (4) 348.4 J K<sup>-1</sup> mol<sup>-1</sup>
- 24. Which of the following represents a correct step in the mechanism of nitration of benzene with conc. HNO<sub>3</sub> / conc. H<sub>2</sub>SO<sub>4</sub>?

$$(1) \bigcirc \stackrel{\text{J}}{\bigcirc} \stackrel{\text{J}}{\longrightarrow} \stackrel{\text{J}}{\bigcirc} \stackrel{\text{H}}{\longrightarrow} \text{NO}_2$$

$$(2) \quad \bigcirc^{\uparrow}_{H}^{NO_{2}} \quad \longrightarrow \quad \bigcirc^{\downarrow}_{H}^{NO_{2}}$$

$$(3) \bigcirc^{NO_2} \longrightarrow \bigcirc^{H}_+ NO_2$$

$$(4) \qquad \stackrel{\text{HSO}_{4}^{2}}{\longrightarrow} \qquad \stackrel{\text{NO}_{2}}{\longrightarrow} \qquad + \quad \text{H}_{2}\text{SO}_{4}$$

$$(5) \begin{array}{c} + NO_2 \\ + H_2SO_4 \end{array}$$

**25**.

CH<sub>2</sub>MgBr

Y

In the reaction sequence given above, the structures of X and Y respectively are,

$$\begin{array}{ccc} \text{(1)} & & & & \text{CO}_2\text{H} \\ & & & \text{CH}_2\text{CH}_2\text{CHCH}_3 \\ & & & \text{OH} \end{array}$$

$$\begin{array}{c} \begin{array}{c} \text{CO}_2\text{MgBr} \\ \\ \text{CH}_2\text{CH}_2\text{CHCH}_3 \\ \\ \text{OMgBr} \end{array}$$

(2) 
$$CH_2OH$$
  
 $CH_2CH_2COCH_3$ 

$$\begin{array}{c} \text{CH}_2\text{OMgBr} \\ \text{CH}_2\text{CH}_3 \\ \text{CH}_2\text{CH}_2 - \text{C} - \text{CH}_3 \\ \text{OMgBr} \end{array}$$

$$(4) \quad \bigcirc \stackrel{\mathrm{CO_2H}}{\bigcirc CH_2\mathrm{CH_2CHCH_3}}$$
 
$$OH$$


- 26. When  $(NH_4)_2CO_3(s)$ ,  $(NH_4)_2Cr_2O_7(s)$  and  $NH_4NO_3(s)$  are heated, the nitrogen containing compounds obtained are respectively,
  - (1)  $NH_3$ ,  $N_2$  and  $NO_2$
- (2) N<sub>2</sub>O, N<sub>2</sub> and NH<sub>3</sub>
   (5) N<sub>2</sub>, NH<sub>3</sub> and N<sub>2</sub>O
- (3)  $NH_3$ ,  $N_2$  and  $N_2O$

- (4)  $N_2$ ,  $N_2$ O and  $NH_3$
- 27. Which of the following would occur soon after connecting a rod of Zn and a rod of Ag immersed in a saturated solution of AgCl and AgCl(s) kept in a beaker as shown in the diagram, by a conductor?

$$Zn^{2+}(aq) + e \longrightarrow Zn(s) E^{o} = -0.76 V$$

 $Ag^{+}(aq) + e \longrightarrow Ag(s) E^{\circ} = 0.80 V$ 

- (1) Zn dissolves, Ag deposits, AgCl(s) dissolves.
- (2) Zn dissolves, Ag dissolves, AgCl(s) dissolves.
- (3) Zn dissolves, Ag dissolves, AgCl(s) deposits.
- (4) Zn deposits, Ag dissolves, AgCl(s) dissolves.
- (5) Chloride concentration in the solution decreases.



28. In the reaction sequence given below, the structures of P and Q respectively are,

$$C_6H_5C\equiv CH \xrightarrow{Hg^{2+}/dil. H_2SO_4} P \xrightarrow{Zn/Hg} Q$$

- (5)  $C_6H_5C=CH_2$  ,  $C_6H_5CHCH_3$  OH OH
- 29. Which of the following statements is incorrect regarding polymers?
  - (1) Bakelite is a thermosetting polymer.
  - (2) Teflon is a thermoplastic polymer.
  - (3) Nylon 6,6 is formed by addition polymerisation between 1,6-diaminohexane and hexanedioic
  - (4) Terelene is formed by condensation polymerisation between ethylene glycol and terephthalic acid.
  - (5) Natural rubber consists of cis-polyisoprene chains.
- 30. An experiment was carried out to find the order (m) with respect to  $S_2O_3^{2-}$  of the reaction  $S_2O_3^{2-}(aq) + 2H^+(aq) \longrightarrow H_2O(l) + SO_2(g) + S(s)$ . Initial rate of the reaction (R) was measured by adding different volumes (v) of 0.01 mol dm<sup>-3</sup>  $S_2O_3^{2-}$  into a solution of an acid. The H<sup>+</sup> concentration of the reaction mixture was kept constant, but the total volume (V) was allowed to vary. Which of the following relations regarding the initial rate of the reaction is correct?
  - (1)  $R \propto \left(\frac{v}{V}\right)^m$  (2)  $R \propto v^m$  (3)  $R \propto v^{\frac{1}{m}}$  (4)  $R \propto \left(\frac{v}{V}\right)^{\frac{1}{m}}$  (5)  $R \propto V^m$
- For each of the questions 31 to 40, one or more responses out of the four responses (a), (b), (c) and (d) given is/are correct. Select the correct response/responses. In accordance with the instructions given on your answer sheet, mark
  - (1) if only (a) and (b) are correct.
  - (2) if only (b) and (c) are correct.
  - (3) if only (c) and (d) are correct.
  - (4) if only (d) and (a) are correct.
  - if any other number or combination of responses is correct.

## Summary of above Instructions

(1)	(2)	(3)	(4)	(5)
Only (a) and (b) are correct	Only (b) and (c) are correct	Only (c) and (d) are correct	Only (d) and (a) are correct	Any other number or combination of responses is correct

- 31. Consider a titration between a weak acid (fixed volume) and a strong base. Which of the following is/are independent of the weak acid concentration?
  - (a) pH at the equivalence point.
  - (b) Volume of the strong base required to reach the end point.
  - (c) Dissociation constant of the weak acid.
  - (d) Value of  $[H^+] \times [OH^-]$  of the solution in the titration flask.

32. Which of the following statements is/are true regarding the molecule given below?

$$\mathbf{CH_3} - \mathbf{C} \equiv \mathbf{C} - \mathbf{CHO}$$

- (a) All four carbon atoms lie in the same plane.
- (b) The angle between  $C_d-H$  and the  $C_d-C_c$  bonds is approximately 120°.
- (c) Between  $C_b$  and  $C_c$ , there are two  $\sigma$ -bonds and one  $\pi$ -bond.
- (d) Between  $C_b$  and  $C_c$ , there is one  $\sigma$ -bond and two  $\pi$ -bonds.
- 33. Which of the following statement/s is/are true with regard to the manufacture of Na<sub>2</sub>CO<sub>3</sub>?
  - (a) CO<sub>2</sub> is one of the raw materials used.
  - (b) The reaction between CO<sub>2</sub> and aqueous NaCl saturated with NH<sub>3</sub> is endothermic.
  - (c) The manufacturing process involves five stages.
  - (d) Most of the NH<sub>3</sub> used in the process can be recovered.
- 34. Temperature must be maintained at a constant value during the experimental determination of the order of an elementary reaction, because,
  - (a) the order of the reaction depends on temperature.
  - (b) the activation energy changes with temperature.
  - (c) the mechanism of the reaction changes with temperature.
  - (d) the rate constant changes with temperature.
- 35. Which of the following statement/s is/are true regarding ethene and ethyne?
  - (a) CaC<sub>2</sub> reacts with water to form ethyne.
  - (b) CaC<sub>2</sub> reacts with water to form ethene.
  - (c) Ethene reacts with ammoniacal AgNO<sub>3</sub> to give a precipitate.
  - (d) Ethyne reacts with ammoniacal Cu<sub>2</sub>Cl<sub>2</sub> to give a precipitate.
- 36. Which of the following statement/s is/are true with regard to halogens?
  - (a) The boiling points of halogens increase down the group.
  - (b) Unlike other halogens, fluorine always has an oxidation state of (-1) except in F<sub>2</sub>.
  - (c) All halogens are good reducing agents.
  - (d) Although fluorine is the most reactive of all the elements in the Periodic Table, it does not react with inert gases.
- 37. For the reaction  $C(s) + CO_2(g) \rightleftharpoons 2 CO(g)$  occurring in a closed rigid container, percentage yields of CO(g) at 700 °C and 800 °C are 60% and 80% respectively. Which of the following statement/s is/are **correct** regarding the above reaction?
  - (a) The reaction is endothermic.
  - (b) The reaction is exothermic.
  - (c) Reverse reaction is favoured by decreasing the temperature.
  - (d) Equilibrium can be shifted towards the reactants by removing C(s).
- 38. Cyclopropane → propene is an elementary reaction.

Which of the following statement/s is/are correct regarding the above reaction?

- (a) Half life of the reaction depends on cyclopropane concentration.
- (b) Rate of the reaction does not depend on propene concentration.
- (c) The fraction of cyclopropane molecules having energy greater than the activation energy increases with increasing temperature.
- (d) Reaction occurs via a bimolecular collision (molecularity = 2).
- 39. Which of the following statement/s is/are true regarding 3-hexene?
  - (a) Does not show geometric isomerism.
  - (b) Shows optical isomerism.
  - (c) The compound obtained when reacted with  $H_2/Pd$  does not show optical isomerism.
  - (d) The compound obtained when reacted with HBr shows optical isomerism.

- 40. Which of the following statements is/are correct with regard to the nitrogen cycle?
  - (a) N<sub>2</sub> in the atmosphere is fixed only by atmospheric and industrial fixation.
  - (b)  $N_2$  is reduced during atmospheric fixation.
  - (c)  $N_2$  is oxidized during industrial fixation.
  - (d) Nitrates and nitrites formed during atmospheric fixation are utilized by plants to make proteins when the rainfall deposit them on the ground.
- In question Nos. 41 to 50, two statements are given in respect of each question. From the Table given below, select the response out of the responses (1), (2), (3), (4) and (5) that best fits the two statements and mark appropriately on your answer sheet.


Response	First Statement	Second Statement
(1) (2) (3) (4) (5)	True True True False False	True, and correctly explains the first statement True, but does <b>not</b> explain the first statement correctly False True False

	First Statement	Second statement
41.	BaCO <sub>3</sub> is more thermally stable than MgCO <sub>3</sub> .	Polarizing power of group two cations decreases down the group.
42.	The lone pair of electrons on nitrogen in an amine has a lower tendency to form a bond with H <sup>+</sup> , than the lone pair of electrons on oxygen in an alcohol.	Nitrogen is less electronegative than oxygen.
43.	A reaction at equilibrium can be driven forward (i.e. shift of equilibrium point to the right) by adding a catalyst.	The catalyst provides a pathway with a low activation energy only to the forward reaction.
44.	$CO_3^{2-}$ and $SO_3^{2-}$ ions have similar shapes.	Central atoms of both $CO_3^{2-}$ and $SO_3^{2-}$ have lone pairs of electrons.
45.	The boiling point of CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> OH is higher than the boiling points of CH <sub>3</sub> CH <sub>2</sub> CHO and CH <sub>3</sub> COCH <sub>3</sub> .	The carbon oxygen double bond is stronger than the carbon oxygen single bond.
46.	A reaction occurring spontaneously in an <b>isolated</b> system <b>always</b> has a negative Gibbs energy change.	A process in an <b>isolated</b> system cannot be changed from outside.
47.	Commonly used soap contain the sodium or potassium salts of fatty acids formed by the reaction of NaOH or KOH with oils and fats.	The reaction of an ester with aqueous NaOH or KOH gives the sodium or potassium salt of the carboxylic acid and the alcohol.
48.	${\rm C_6H_5Br}$ does not react easily with NaOH to form ${\rm C_6H_5OH}$ .	The phenyl carbocation is very stable.
49.	When an aqueous solution of a weak acid is diluted, both the fraction of dissociated acid molecules and pH of the medium are increased.	Dissociation of weak acid molecules occur in such a way that the acid dissociation constant $K_{\rm a}$ remains constant.
50.	In the presence of sunlight CO <sub>2</sub> is fixed in green plants.	Increase of CO <sub>2</sub> level in the atmosphere cannot be controlled by green plants.

# The Periodic Table

	1																	2
1	H		_															He
	3	4											5	6	7	8	9	10
2	Li_	Be											В	C	N	O	F	Ne
	11	12											13	14	15	16	17	18
3	Na	Mg											Al	Si	P	S	CI	Ar
	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
4	K	Ca	Sc	_Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zń	Ga	Ge	As	Se	Br	Kr
	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
	55	56	La-	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
6	Cs	Ba	Lu	Hf	Ta	$\mathbf{w}$	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
	87	88	Ac-	104	105	106	107	108	109	110	111	112	113					
7	Fr	Ra	Lr	Rf	Db	Sg	Bh	Hs	Mt	Uun	Uuu	Uub	Uut					

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr



ପିପତ୍ର ଡ ଷିଡିରଡି ଫ୍ରିମିମ୍ମି /( $\wp$ ୍ରଡ଼ । ଧର୍ଣ୍ଡାଧ୍ୟ ଧର୍ଣ୍ଡାଧ୍ୟ ପ୍ରଥମ ନଥିଲେ Reserved]

ලී ලංකා විභාග දෙපාර්තමේත්තුව ලී ලංකා විභාග දෙපාර්තමේ**නි ලියිකා මිහිත පෙපාර්තමේන්තුව**ා විභාග දෙපාර්තමේන්තුව ලී ලංකා විභාග දෙපාර්තමේන්තුව මූහත්කතට பුර්ධනවල් නිකාක්තයගත්ව මූහත්කතට පුද්ධන්දී නිකාක්තයගේ මුහත්කතට පුර්ධන්දී නිකාක්තයගේ මුහත්කතට ප්රධානවල් නිකාක්තයගේ Department of Examinations, Sri Lanka Department of මහත්තය ප්රධාන ප්රධාන විභාග දෙපාර්තමේන්තුව ලී ලංකා විභාග දෙපාර්තමේන්තුව මුහත්කතට ප්රධානවේ නිකාක්තයගේ මුහත්කතට ප්රධානවේ නිකාක්තය ප්රධානවේ නිකාක්තය සහ මුහත්කතට ප්රධානවේ නිකාක්තය සහ මුහත්කතට ප්රධානවේ නිකාක්තය සහ මුහත්කතට ප්රධානවේ නිකාක්තිකතට ප්රධානවේ නිකාක්තිකතට ප්රධානවේ නිකාක්තිකතට ප්රධානවේ නිකාක්තිකතට ප්රධානවේ නිකාක්තිකතට ප්රධානවේ නිකාක්තය සහ ප්රධානවේ නිකාක්තිකතට ප්රධානවේ නිකාක්තිකතට ප්රධානවේ නිකාක්තිකට ප්රධානවේ නිකාක්තිකතට ප්රධානවේ නිකාක්තිකතට ප්රධානවේ නිකාක්තිකට ප්රධානවේ නිකාක්තිකට ප්රධානවේ නිකාක්තිකට ප්රධානවේ නිකාක්තිකට ප්රධානවේ සහ ප්රධාන ප්

අ<del>பங்கள் பேற்ற கல்கின் பற (උසස් பேத) විභාගය, 2018 අගෝස්තු</del> கல்விப் பொதுத் தராதரப் பத்திர (உயர் தர)ப் பரீட்சை, 2018 ஓகஸ்ற் General Certificate of Education (Adv. Level) Examination, August 2018

රසායන විදාහාව II இரசாயனவியல் II Chemistry II

02 E II

17.08.2018 / 0830 - 1140

*පැය තුනයි* மூன்று மணித்தியாலம் **Three hours** 

අමතර කියවීම් කාලය - මිනිත්තු 10 යි மேலதிக வாசிப்பு நேரம் - 10 நிமிடங்கள் Additional Reading Time - 10 minutes

Use additional reading time to go through the question paper, select the questions and decide on the questions that you give priority in answering.

Index No.: .....

- \* A Periodic Table is provided on page 16.
- \* Use of calculators is not allowed.
- \* Universal gas constant,  $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
- \* Avogadro constant,  $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$
- \* In answering this paper, you may represent alkyl groups in a condensed manner.

Example: H—C——C— group may be shown as CH<sub>3</sub>CH<sub>2</sub>—
H H

## □ PART A — Structured Essay (pages 2 - 8)

- \* Answer all the questions on the question paper itself.
- \* Write your answer in the space provided for each question. Please note that the space provided is sufficient for the answer and that extensive answers are not expected.

#### $\Box$ PART B and PART C — Essay (pages 9 - 15)

- \* Answer four questions selecting two questions from each part. Use the papers supplied for this purpose.
- \* At the end of the time allotted for this paper, tie the answers to the three Parts A, B and C together so that Part A is on top and hand them over to the Supervisor.
- \* You are permitted to remove only Parts B and C of the question paper from the Examination Hall.

#### For Examiner's Use Only

Part	Question No.	Marks				
	1					
A	2					
	3					
	4					
	5					
В	6					
	7					
	8					
C	9					
	10					
Total						
Percentage						

#### Final Mark

In Numbers	
In Letters	

#### **Code Numbers**

Marking Examiner 1	
Marking Examiner 2	
Checked by :	
Supervised by:	

(v) Consider the hydrides of the elements in the group to which X belongs, which are analogous to Y. Sketch the variation in boiling points of these hydrides (including Y) in the graph below. In your sketch indicate the hydrides using their chemical formulae. (Note: Values of boiling points are not required.)
Boiling point A
(vi) Give reasons for the variation in boiling points in part (v) above.
<ul> <li>(vii) I. Write what you would observe when an excess of an aqueous solution of Y is added to a solution of Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>.</li> <li>II. Write the chemical formula of the species that gives rise to your observation in part I above.</li> </ul>
(viii) Give one chemical test to identify Y.
Test:
Observation:
(ix) $\mathbf{Z}$ is an oxo-acid of $\mathbf{X}$ and a strong oxidizing agent.
I. Identify Z.
II. State the products obtained when hot concentrated Z reacts with sulphur.
(b) A and B are compounds of two p-block elements that belong to the same group in the Periodic Table. A exists as a colourless, odourless liquid at room temperature and atmospheric pressure. It is also found in the gaseous and solid states. The solid state of A is less dense than its liquid state. Ionic and polar compounds are readily soluble in A.  B is a colourless gas at room temperature and atmospheric pressure. A filter paper moistened with lead acetate turns black on treatment with B.  (i) Identify A and B.
$A = \dots B = \dots$
(ii) Sketch the shapes of A and B showing lone pairs of electrons where necessary.

Do not write in this column.

47

0

` '	Giving reasons, state	wnether	01 1		110 1415	er bond angle.	
			• • • • • • • • • • • • • • • • • • • •			.,,,,,	
			•••••				,
(iv)	In each of the follo action of A.	wing inst	ances, g	ive a b	alanceo	d chemical equation to in	dicate the
	I. A as an acid:						
	II. A as a base:						
(v)	Write the balanced of	hemical	equation	for the	e reacti	on of <b>B</b> with aqueous lea	d acetate.
			• • • • • • • • • •		• • • • • • • • • • • • • • • • • • • •		
(vi)	I. Write what you solution of BiCl <sub>3</sub>		serve wh	nen A. a	and <b>B</b> a	are added <b>separately</b> to an	acidified
	with A (excess)	:				with $B:$	
	II. Write balanced of	hemical	equations	s for y	our obs	servations in part I above	•
						•••••	
			•••••	• • • • • • • • •	• • • • • • • • •	•••••	(4.0 marks)
		-	-		-	vas carried out at 25 °C. In 1.10 mol of <b>B</b> in distilled w	-
eaction olume	mixture was made by	dissolving dissolving the c	ng 0.10 n	nol of A	A and 0	vas carried out at 25 °C. In 1.10 mol of <b>B</b> in distilled w is solution with time is sho	ater (total
eaction olume	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (mo	dissolving in the column distance of the colu	ng 0.10 n	nol of A	A and 0	.10 mol of B in distilled w	ater (total
eaction olume	mixture was made by 100.00 cm <sup>3</sup> ). Variatio	dissolving in the column distance of the colu	ng 0.10 n	nol of A	A and 0 A in th	.10 mol of <b>B</b> in distilled wis solution with time is sho	ater (total
eaction olume	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (mo	dissolving in the column distance of the colu	ng 0.10 n	nol of A	A and 0 A in th	.10 mol of B in distilled w	ater (total
eaction olume	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (mo	dissolvin n in the c	ng 0.10 n	nol of A	A and 0 A in th	.10 mol of <b>B</b> in distilled wis solution with time is sho	ater (total
eaction olume	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (mo	dissolving in the column distance of the colu	ng 0.10 n	nol of A	A and 0 A in th	.10 mol of <b>B</b> in distilled wis solution with time is sho	ater (total
eaction olume	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (mo	dissolvin n in the c	ng 0.10 n	nol of A	A and 0 A in th	.10 mol of <b>B</b> in distilled wis solution with time is sho	ater (total
eaction olume	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (mo	dissolvir n in the c	ng 0.10 n	nol of A	A and 0 A in th	.10 mol of <b>B</b> in distilled wis solution with time is sho	ater (total
eaction olume	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (mo	dissolvir n in the c	ng 0.10 n	nol of A	A and 0 A in th	.10 mol of <b>B</b> in distilled wis solution with time is sho	ater (total
eaction rolume graph.	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (model)  1.0  0.5	dissolvin n in the cold dm <sup>-3</sup> )	ng 0.10 n oncentral	nol of Ation of	A and 0 A in th	.10 mol of <b>B</b> in distilled was solution with time is shown that the control of t	rater (total own in the
eaction rolume graph.	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (model)  1.0  0.5	dissolvin n in the cold dm <sup>-3</sup> )	ng 0.10 n oncentral	nol of Ation of	A and 0 A in th	.10 mol of <b>B</b> in distilled was solution with time is shown that the solution with t	rater (total own in the
eaction volume graph.	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (model)  1.0  0.5	dissolvin n in the cold dm <sup>-3</sup> )	ng 0.10 n oncentral	nol of Ation of	A and 0 A in th	.10 mol of <b>B</b> in distilled was solution with time is shown that the solution with t	rater (total own in the
eaction volume graph.	n mixture was made by 100.00 cm <sup>3</sup> ). Variatio concentration (model)  1.0  0.5	dissolvin n in the cold dm <sup>-3</sup> )	ng 0.10 n oncentral	nol of Ation of	A and 0 A in th	.10 mol of <b>B</b> in distilled was solution with time is shown that the solution with t	rater (total own in the
eaction volume graph.  i) Calconnection Calc	n mixture was made by 100.00 cm³). Variatio concentration (mo 1.0 0.5 0.5	dissolving in the cool dm <sup>-3</sup> )  2.0  A (in motorward re-	4.0 oles) reaction be	6.0 cted du	A and 0 A in th	.10 mol of <b>B</b> in distilled was solution with time is shown that the solution with t	rater (total own in the reaction.
eaction volume graph.  i) Calconnection Calc	n mixture was made by 100.00 cm³). Variatio concentration (mo 1.0 0.5 0.5 0.0 culate the amount of	2.0 A (in mo	4.0 oles) reaction be	6.0 cted du	A and 0 A in the	[A]  [A]  time (min.)  e first 4.0 minutes of the	rater (total own in the reaction.

(c) (i) Give the mechanism of the following reaction.

$$C_2H_5OH + HBr \longrightarrow C_2H_5Br + H_2O$$

- (ii) State whether the above reaction is a nucleophilic substitution reaction or an electrophilic substitution reaction. Identify the nucleophile or electrophile as appropriate.
- (iii) State giving reasons which of the two compounds, phenol ( $C_6H_5OH$ ) or ethanol ( $C_2H_5OH$ ) is more acidic.

(3.0 marks)

### PART C - ESSAY

Answer two questions only. (Each question carries 15 marks.)

**8.** (a) An aqueous solution **P** contains **two** cations and **two** anions. The following experiments were carried out to identify these cations and anions.

### Cations

	Experiment	Observation
	${\bf P}$ was acidified with dilute HCl and ${\bf H}_2{\bf S}$ was bubbled through the solution.	A clear solution was obtained.
2	The above solution was boiled till all the $\rm H_2S$ was removed. A few drops of conc. $\rm HNO_3$ were added and the solution was heated further. The resulting solution was cooled and $\rm NH_4Cl/NH_4OH$ was added.	A brown precipitate (Q) was formed.
	Q was removed by filtration and H <sub>2</sub> S was bubbled through the filtrate.	A pale pink precipitate (R) was formed.
	$\bf R$ was removed by filtration and the filtrate was boiled till all the $\rm H_2S$ was removed. $\rm (NH_4)_2CO_3$ was added to the solution.	A clear solution was obtained.
<b>⑤</b>	Dilute NaOH was added to a fresh portion of P.	A dirty-green precipitate and a while precipitate were formed.

## Experiments for precipitates Q and R:

	Experiment	Observation						
6	${f Q}$ was dissolved in dil. ${\ensuremath{HNO}}_3$ and a salicylic acid solution							
	was added.	obtained.						
	${f R}$ was dissolved in dilute acid and dil. NaOH was added to the solution.	A white precipitate was formed. It turned brown on standing.						

#### **Anions**

		Test	Observation					
8	I. BaCl <sub>2</sub> solu	tion was added to P.	A white precipitate was formed.					
		precipitate was separated by filtration and as added to the precipitate.	The white precipitate was not dissolved.					
9		and chloroform were added to a portion of from ® II, and the mixture was throughly						

- (i) Identify the two cations and the two anions in solution P. (Reasons are not required.)
- (ii) Write the chemical formulae of the precipitates Q and R.
- (iii) Give reasons for the following:
  - I. Removal of H<sub>2</sub>S in experiment ② for cations.
  - II. Heating with conc. HNO<sub>3</sub> in experiment ② for cations.

(7.5 marks)

(b) The sample X contains lead, copper and an inert material. The following procedure was carried out to analyse lead and copper in X.

#### Procedure:

A mass of 0.285 g of X was dissolved in a slight excess of dil.HNO<sub>3</sub>. A clear solution was obtained. A NaCl solution was added to the resulting clear solution. A white precipitate (Y) was formed. The precipitate was separated by filtration and the precipitate (Y) and filtrate (Z) were analysed separately.

## Precipitate (Y)

The precipitate was dissolved in hot water. A solution of  $K_2CrO_4$  was added in excess. A yellow precipitate was formed. The precipitate was separated by filtration and dissolved in dil. HNO<sub>3</sub>. An orange coloured solution was obtained. Excess KI was added to this solution and the liberated  $I_2$  was titrated with 0.100 mol dm<sup>-3</sup> Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, with starch as the indicator. The volume of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> required to reach the end point was 27.00 cm<sup>3</sup>. (Assume that the NO<sub>3</sub> ions do **not** interfere with the titration.)

## Filtrate (Z)

The filtrate was neutralized and excess KI was added to it. The liberated  $I_2$  was titrated with 0.100 mol dm<sup>-3</sup> Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, with starch as the indicator. The volume of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> required to reach the end point was 15.00 cm<sup>3</sup>.

(Note: Assume that the inert material was soluble in dil. HNO<sub>3</sub> and did not interfere with the experiment.)

- (i) Calculate the mass percentages of lead and copper in X. Write balanced chemical equations where relevant.
- (ii) What is the colour change at the end point in the titration carried out in the analysis of precipitate Y?

$$(Cu = 63.5, Pb = 207)$$

(7.5 marks)


- 9. (a) The following questions are based on the environment and related issues.
  - (i) Identify three greenhouse gases that contribute to global warming. State two consequences of global warming.
  - (ii) Global environmental issues caused by coal power plants are well known. Identify **one** such issue that contributes **significantly** to change in certain water quality parameters in rivers and lakes.
  - (iii) Name the chemical species responsible for the environmental issue identified in (ii) above and state **three** water quality parameters that are likely to be affected by this issue.
  - (iv) Identify **two** environmental issues that change (increase or decrease) the ozone level in the atmosphere and explain briefly how these changes take place with the aid of balanced chemical equations.
  - (v) I. "Most of the harmful gases in vehicle exhausts are converted to relatively harmless gases by catalytic converters." Briefly explain this statement.
    - II. Name the harmful gas (except CO<sub>2</sub>) that is not converted to a less harmful gas by the catalytic converter. State briefly how this harmful gas is formed in the vehicle engine.

(7.5 marks)

(b) The flow chart given below shows the production of two important compounds  $P_1$  and  $P_2$  and three other important compounds  $P_3$ ,  $P_4$  and  $P_5$  derived from them.  $P_1$  is used as a raw material in the manufacture of  $Na_2CO_3$ .  $P_3$  can be manufactured by the reaction between  $P_1$  and  $P_2$ .  $P_3$  is used as a fertilizer and as an explosive.  $P_1$  is also used in the manufacture of  $P_4$  which is a widely used fertilizer.  $P_4$  is used to synthesize an important thermosetting polymer  $P_5$ .

M Manufacturing process
PC Physical/chemical process to obtain raw material
P Product
S Source of raw material

X Unreacted raw material(s) / substance discharged to the atmosphere during physical and/or chemical process



Answer the following questions based on the above flow chart.

- (i) Identify  $P_1$ ,  $P_2$ ,  $P_3$ ,  $P_4$  and  $P_5$
- (ii) Identify  $\mathbf{R}_1$ ,  $\mathbf{R}_2$  and  $\mathbf{R}_3$
- (iii) Identify  $X_1$ ,  $X_2$  and  $X_3$
- (iv) Identify S.
- (v) Briefly state the processes taking place in  $\mathbf{PC}_1$  and  $\mathbf{PC}_2$  giving balanced chemical equations where applicable.
- (vi) Identify manufacturing processes  $M_1$ ,  $M_2$  and  $M_3$ . (e.g. contact process or manufacture of  $H_2SO_4$ .)
- (vii) Give balanced chemical equations with appropriate conditions, for reactions taking place in  $M_1$ ,  $M_2$  and  $M_3$ .
- (viii) I. Give one use of each compound  $\mathbf{P_1}$  and  $\mathbf{P_2}$  other than those mentioned above.
  - II. Give one use of  $\mathbf{R}_1$  in the manufacturing process  $\mathbf{P}_1$  other than being used as a raw material. (7.5 marks)

10.(a) A and B are complex ions, (i.e. metal ion and ligands coordinated to it) with an octahedral geometry. They have the same atomic composition of MnC<sub>5</sub>H<sub>3</sub>N<sub>6</sub>. In each complex ion, two types of ligands are coordinated to the metal ion. When an aqueous solution containing A is treated with a potassium salt, the coordination compound C is formed. C gives four ions in aqueous solution. When an aqueous solution containing B is treated with a potassium salt the coordination compound D is formed. D gives three ions in aqueous solution. Both C and D have an octahedral geometry.

(Note: The oxidation states of manganese in A and B do not change on treatment with the potassium salt).

- (i) Identify the ligands coordinated to manganese in A and B.
- (ii) Give the structures of A, B, C and D.
- (iii) Write the electronic configurations of the manganese ions in A and B.
- (iv) Write the IUPAC names of C and D.

(7.5 marks)


- (b) (i) I. Write the reduction half reaction corresponding to the electrode, Ag(s) | AgCl(s) | Cl<sup>-</sup>(aq).
  - II. State whether the electrode potential of Ag(s) | AgCl(s) | Cl<sup>-</sup>(aq) depends on the Ag<sup>+</sup> concentration in the solution. Explain your answer.
  - (ii) Consider the following reaction.

$$Fe(s) + 2H^{+}(aq) + \frac{1}{2}O_{2}(g) \longrightarrow Fe^{2+}(aq) + H_{2}O(l)$$

- I. Write the oxidation and reduction half reactions relevant to the above reaction.
- II. Given that the above reaction is the cell reaction of an electrochemical cell, determine the standard electromotive force of the cell.

$$E_{Fe^{2+}(aq)/Fe(s)}^{o} = -0.44 \text{ V}$$
  $E_{H^{+}(aq)/O_{2}(g)/H_{2}O(l)}^{o} = 1.23 \text{ V}$ 

(iii) A constant current of 100 mA was passed through 100.00 cm<sup>3</sup> of a 0.10 mol dm<sup>-3</sup> aqueous CaBr<sub>2</sub> solution as shown in the diagram. The temperature of the system was maintained at 25 °C.



- I. Write the oxidation and reduction reactions that take place at the electrodes.
- II. Calculate the time taken for the commencement of precipitation of  $Ca(OH)_2(s)$ . Solubility product of  $Ca(OH)_2$  at 25 °C is  $1.0 \times 10^{-5}$  mol<sup>3</sup> dm<sup>-9</sup>. Neglect the ionization of water. Assume that the volume of the aqueous phase remains constant.

(7.5 marks)

# The Periodic Table

		-																
	1																	$\lceil 2 \rceil$
1	H		,															He
	3	4											5	6	7	8	9	10
2	Li	Be											В	C	N	O	F	Ne
	11	12											13	14	15	16	17	18
3	Na	Mg											AI	Si	P	S	Cl	Ar
	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
5	Rb	Sr	Y_	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Те	1	Xe
	55	56	La-	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
6	Cs	Ba	Lu	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	TI	Pb	Bi	Po	At	Rn
	87	88	Ac-	104	105	106	107	108	109	110	111	112	113					I
7	Fr	Ra	Lr	Rf	Db	Sg	Bh	Hs	Mt	Uun	Uuu	Uub	Uut					

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr



විභාග ඉලක්ක පහසුවෙන් ජයගන්න

පසුගිය විභාග පුශ්න පතු



 Past Papers
 Model Papers
 Resource Books for G.C.E O/L and A/L Exams





້ວິສາທ ຈູ@ສໄສ ປ໌ຜທສໄສ Knowledge Bank











Website WWW.Iol.Ik



071 777 4440